





Gestational Diabetes PowerPlan – Ordering and administration

Ordering the Gestation Diabetes Powerplan


1. Open the *Orders* page and click *add*.
2. Search for the *Gestation Diabetes – Maternity PowerPlan*.




3. Choose the relevant phase/s of the PowerPlan from the side *View* menu.



4. Complete the relevant *Patient Care* tasks and add Medications (if applicable) to the relevant phase by clicking *add to phase*.



5. If subcutaneous insulin is required to be ordered, this needs to be completed using the relevant *Subcutaneous Insulin Careset* or *Supplemental Insulin PowerPlan* after initiating the Gestational Diabetes PowerPlan (Caresets and PowerPlans cannot be added to a PowerPlan phase).




6. When all relevant orders have been added, click *Initiate Now* and then *Sign*.
7. Refresh the *Orders* screen and check the *MAR* to ensure any medication orders are correct.

Administering orders from the Gestational Diabetes PowerPlan

- Patient care and medication tasks ordered as part of the Gestational Diabetes PowerPlan will appear in *CareCompass* and the *MAR*.

CareCompass:



MAR:

Medications	25-Apr-2019 12:00 AEST	25-Apr-2019 8:00 AEST	25-Apr-2019 7:30 AEST	24-Apr-2019 20:00 AEST	24-Apr-2019 18:00 AEST	24-Apr-2019 17:00 AEST
gentamicin						8 unit(s) Not given within 5 days.
insulin novorapid (novorapid FlexPen) 0-0-8-0 unit(s) W, Pen, Subcutaneous, start: 24-Apr-2019 17:00:00, stop: 25-Apr-2019 16:59:00, (ultra short acting) for Wednesday (with start of meal), Indication: Diabetes						
insulin novorapid Blood glucose, Capillary						
insulin novorapid (novorapid FlexPen) 4-6-8-0 unit(s) Th, Pen, Subcutaneous, start: 25-Apr-2019 08:00:00, stop: 26-Apr-2019 07:59:00, (ultra short acting) for Thursday (with start of meal), Indication: Diabetes	6 unit(s) Not given within 5 days.	4 unit(s) Not given within 5 days.				
insulin novorapid Blood glucose, Capillary						
insulin subcutaneous dose check Adult 1 EA, Other-enter comment, THREE times a day (before food), start: 24-Apr-2019 17:00:00, Insulin subcutaneous injection has been ordered for this patient, please ensure doses have been ordered., Indication: insulin subc...	1 EA Not given within 5 days.			1 EA Not given within 5 days.		1 EA Not given within 5 days.
insulin subcutaneous dose check Adult						
metformin (metformin 500 mg oral IMMEDI... 500 mg = 1 tab(s), Tablet, Oral, TWICE a day (with or after food), start: 24-Apr-2019 18:00:00, Indication: Gestational Diabetes		500 mg Not given within 5 days.			500 mg Not given within 5 days.	

- For further information on how to administer regular and supplemental insulin, please refer the *Regular and Supplemental Insulin: Administration* Quick Reference Guide.